

Quantum Mechanics in a Time-Asymmetric Universe: On the Nature of the Initial Quantum State

Eddy Keming Chen

In a quantum universe with a strong arrow of time, we postulate a low-entropy boundary condition (the past hypothesis, PH) to account for the temporal asymmetry. In this article, I show that the PH also contains enough information to simplify the quantum ontology and define a natural initial condition. First, I introduce ‘density-matrix realism’, the thesis that the quantum state of the universe is objective and impure. This stands in sharp contrast to wave-function realism, the thesis that the quantum state of the universe is objective and pure. Second, I suggest that the PH is sufficient to determine a natural density matrix, which is simple and unique. This is achieved by what I call the ‘initial projection hypothesis’: the initial density matrix of the universe is the (normalized) projection onto the PH-subspace (in the Hilbert space). Third, because the initial quantum state is unique and simple, we have a strong case for the nomological thesis: the initial quantum state of the universe is on a par with laws of nature. This new package of ideas has several interesting implications, including on the harmony between statistical mechanics and quantum mechanics, theoretical unity of the universe and the subsystems, and the alleged conflict between Humean supervenience and quantum entanglement.

1. Introduction

In the foundations of quantum mechanics, it has been argued that the wave function (pure state) of the universe represents something objective and not something merely epistemic. Let us call this view ‘wave-function realism’. There are many realist proposals for how to understand the wave function. Some argue that it represents things in the ontology, either a physical field propagating on a fundamental high-dimensional space, or a multi-field propagating on the three-dimensional physical space. Others argue that it is in the ‘nomology’—having the same status as laws of nature. Still others argue that it might belong to a new ontological category.¹

¹ See (Albert [1996]; Loewer [1996]; Wallace and Timpson [2010]; Ney [2012]; North [2013]; Maudlin [2013]; Goldstein and Zanghi [2013]; Miller [2014]; Esfeld [2014]; Bhogal and Perry [2015]; Callender [2015]; Esfeld and Deckert [2017]; Chen [2017], [unpublished-a], [unpublished-b]; Hubert and Romano

Wave-function realism has generated much debate. In fact, it has been rejected by many people, notably by quantum Bayesians, and various anti-realists and instrumentalists. As a scientific realist, I do not find their arguments convincing. In previous papers, I have assumed and defended wave-function realism. Nevertheless, in this article I want to argue for a different perspective, for reasons related to the origin of time-asymmetry in a quantum universe.

To be sure, realism about the universal wave function is highly natural in the context of standard quantum mechanics and various realist quantum theories such as Bohmian mechanics (BM), Ghirardi–Rimini–Weber (GRW) spontaneous collapse theories, and Everettian quantum mechanics (EQM). In those theories, the universal wave function is indispensable to the kinematics and the dynamics of the quantum system. However, as I would like to emphasize in this article, our world is not just quantum-mechanical. We also live in a world with a strong arrow of time (large entropy gradient). There are thermodynamic phenomena that we hope to explain with quantum mechanics and quantum statistical mechanics (QSM). A central theme of this article is to suggest that QSM is highly relevant for assessing the fundamentality and reality of the universal wave function.

We will take a close look at the connections between the foundations of QSM and various solutions to the quantum measurement problem. When we do, we realize that we do not need to postulate a universal wave function. We need only certain ‘coarse-grained’ information about the quantum macro-state, which can be represented by a density matrix. A natural question is: can we understand the universal quantum state as a density matrix rather than a wave function? That is, can we take an ‘ontic’ rather than an ‘epistemic’ attitude towards the density matrix?

The first step of this article is to argue that we can. I call this view ‘density-matrix realism’, the thesis that the actual quantum state of the universe is objective (as opposed to subjective or epistemic) and impure (mixed). This idea may be unfamiliar to some people, as we are used to take the mixed states to represent our epistemic uncertainties of the actual pure state (a wave function). The proposal here is that the density matrix directly represents the actual quantum state of the universe; there is no further fact about which is the actual wave function. In this sense, the density matrix is ‘fundamental’. In fact, this idea has come up in the foundations of physics.² In the first step, we provide a systematic discussion of density-matrix realism by reformulating BM, GRW theories, and EQM in terms of a fundamental density matrix.

The second step is to point out that density-matrix realism allows us to combine quantum ontology with time-asymmetry in a new way. In classical and quantum

[2018]). For a survey on this literature, see (Chen [2019]). Notice that this is not how Albert, Loewer, or Ney characterizes wave-function realism. For them, to be a wave-function realist is to be a realist about the wave function and a fundamental high-dimensional space—the ‘configuration space’. For the purpose of this article, let us use ‘wave-function realism’ to designate just the commitment that the wave function represents something objective.

² See, for example, (Dürr *et al.* [2005]; Maroney [2005]; Wallace [2012], [unpublished], [forthcoming]).

statistical mechanics, thermodynamic time-asymmetry arises from a special boundary condition called the past hypothesis (PH).³ I suggest that the information in the PH is sufficient to determine a natural density matrix. I postulate the ‘initial projection hypothesis’ (IPH): the quantum state of the universe at t_0 is given by the (normalized) projection onto the PH-subspace, which is a particular low-dimensional subspace in the total Hilbert space. The conjunction of this hypothesis with density-matrix realism pins down a unique initial quantum state. Since the IPH is as simple as the PH, we can use arguments for the simplicity of the latter (which is necessary for it to be a law of nature) to argue for the simplicity of the former. We can thus infer that the initial quantum state is very simple.

The third step is to show that, because of the simplicity and the uniqueness of the initial quantum state (now given by a fundamental density matrix), we have a strong case for the nomological thesis: the initial quantum state of the world is on a par with laws of nature. It is a modal thesis. It implies that the initial quantum state of our world is nomologically necessary; it could not have been otherwise.

As we shall see, this package of views has interesting implications for the reduction of statistical-mechanical probabilities to quantum mechanics, the dynamic and kinematic unity of the universe and the subsystems, the nature of the initial quantum state, and Humean supervenience (HS) in a quantum world.

Here is the roadmap of the article. First, in Section 2, I review the foundations of quantum mechanics and QSM. In Section 3, I introduce the framework of density-matrix realism and provide some illustrations. In Section 4, I propose the IPH in the framework of density-matrix realism. In Section 5, I discuss their implications for statistical mechanics, dynamic unity, and kinematic unity. In Section 6, I suggest that they provide a strong case for the nomological thesis and a new solution to the conflict between quantum entanglement and HS.

2. Foundations of Quantum Mechanics and Statistical Mechanics

In this section, we first review the foundations of quantum mechanics and statistical mechanics. As we shall see in the next section, they suggest an alternative to wave-function realism.

2.1. Quantum mechanics

Standard quantum mechanics is often presented with a set of axioms and rules about measurement. First, there is a quantum state of the system, represented by a wave function ψ . For a spin-less N -particle quantum system in \mathbb{R}^3 , the wave function is a (square-integrable) function from the configuration space \mathbb{R}^{3N} to the complex numbers \mathbb{C} . Second, the wave function evolves in time according to the Schrödinger equation:

³ For an extended discussion, see (Albert [2000]).

$$i\hbar \frac{\partial \psi}{\partial t} = H\psi. \quad (1)$$

Third, the Schrödinger evolution of the wave function is supplemented with collapse rules. The wave function typically evolves into superpositions of macro-states, such as the cat being alive and the cat being dead. This can be represented by wave functions on the configuration space with disjoint macroscopic supports X and Y . During measurements, which are not precisely defined processes in the standard formalism, the wave function undergoes collapses. Moreover, the probability that it collapses into any particular macro-state X is given by the Born rule:

$$P(X) = \int_X |\psi(x)|^2 dx. \quad (2)$$

As such, quantum mechanics is not a candidate for a fundamental physical theory. It has two dynamical laws: the deterministic Schrödinger equation and the stochastic collapse rule. What are the conditions for applying the former, and what are the conditions for applying the latter? Measurements and observations are extremely vague concepts. Take a concrete experimental apparatus for example. When should we treat it as part of the quantum system that evolves linearly and when should we treat it as an ‘observer’, that is, something that stands outside the quantum system and collapses the wave function? That is, in short, the quantum measurement problem.⁴

Various solutions have been proposed regarding the measurement problem. BM solves it by adding particles to the ontology and an additional guidance equation for the particles’ motion. GRW theories postulate a spontaneous collapse mechanism. EQM simply removes the collapse rules from standard quantum mechanics and suggest that there are many (emergent) worlds, corresponding to the branches of the wave function, which are all real. My aim here is not to adjudicate among these theories. Suffice it to say that they are all quantum theories that remove the centrality of observations and observers.

To simplify the discussions, I will use BM as a key example.⁵ In BM, in addition to the wave function that evolves unitarily according to the Schrödinger equation, particles have precise locations, and their configuration $Q = (Q_1, Q_2, \dots, Q_N)$ follows the guidance equation:

$$\frac{dQ_i}{dt} = \frac{\hbar}{m_i} \text{Im} \frac{\nabla_i \psi(q)}{\psi(q)} (q = Q). \quad (3)$$

Moreover, the initial particle distribution is given by the quantum equilibrium distribution:

$$\rho_\omega(q) = |\psi(q, t_0)|^2. \quad (4)$$

⁴ See (Bell [1990]; Myrvold [2017]) for introductions to the quantum measurement problem.

⁵ See (Dürr *et al.* [1992]) for a rigorous presentation of BM and its statistical analysis.

By equivariance, if this condition holds at the initial time, then it holds at all times. Consequently, BM agrees with standard quantum mechanics with respect to the Born rule predictions (which are all there is to the observable predictions of quantum mechanics). For a universe with N particles, let us call the wave function of the universe the ‘universal wave function’ and denote it by $\Psi(q_1, q_2, \dots, q_N)$.

2.2. Quantum statistical mechanics

Statistical mechanics concerns macroscopic systems such as gas in a box. It is an important subject for understanding the arrow of time. For concreteness, let us consider a quantum-mechanical system with N fermions (with $N > 10^{20}$) in a box $\Lambda = [0, L]^3 \subset \mathbb{R}^3$ and a Hamiltonian, \hat{H} . I will first present the ‘individualistic’ view followed by the ‘ensamblist’ view of QSM.⁶ I will include some brief remarks comparing QSM to classical statistical mechanics (CSM), the latter of which may be more familiar to some readers.

- (1) Micro-state: At any time t , the micro-state of the system is given by a normalized (and anti-symmetrized) wave function:

$$\psi(q_1, \dots, q_N) \in \mathcal{H}_{total} = L^2(\mathbb{R}^{3N}, \mathbb{C}^k), \quad \|\psi\|_{L^2} = 1, \quad (5)$$

where $\mathcal{H}_{total} = L^2(\mathbb{R}^{3N}, \mathbb{C}^k)$ is the total Hilbert space of the system. (In CSM, the micro-state is given by the positions and the momenta of all the particles, represented by a point in phase space.)

- (2) Dynamics: The time-dependence of $\psi(q_1, \dots, q_N; t)$ is given by the Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = H\psi. \quad (6)$$

(In CSM, the particles move according to the Hamiltonian equations.)

- (3) Energy shell: The physically relevant part of the total Hilbert space is the subspace (‘the energy shell’):

$$\mathcal{H} \subseteq \mathcal{H}_{total}, \quad \mathcal{H} = \text{span}\{\varphi_\alpha : E_\alpha \in [E, E + \delta E]\}. \quad (7)$$

This is the subspace (of the total Hilbert space) spanned by energy eigenstates φ_α whose eigenvalues E_α belong to the $[E, E + \delta E]$ range. Let $D = \dim \mathcal{H}$, the number of energy levels between E and $E + \delta E$. We only consider wave functions ψ in \mathcal{H} .

- (4) Measure: The measure μ is given by the normalized surface area measure on the unit sphere in the energy subspace $\mathcal{S}(\mathcal{H})$.

⁶ Here I follow the discussions in (Goldstein *et al.* [2010a]; Goldstein and Tumulka [2011]).

- (5) Macro-state: With a choice of macro-variables (suitably ‘rounded’ in the sense of Von Neumann [1955]), the energy shell \mathcal{H} can be orthogonally decomposed into macro-spaces:

$$\mathcal{H} = \oplus_v \mathcal{H}_v, \quad \sum_v \dim \mathcal{H}_v = D. \quad (8)$$

Each \mathcal{H}_v corresponds more or less to small ranges of values of macro-variables that we have chosen in advance. (In CSM, the phase space can be partitioned into sets of phase points. They will be the macro-states.)

- (6) Non-unique correspondence: Typically, a wave function is in a superposition of macro-states and is not entirely in any one of the macrospace. However, we can make sense of situations where ψ is (in the Hilbert space norm) very close to a macro-state \mathcal{H}_v :

$$\langle \psi | P_v | \psi \rangle \approx 1, \quad (9)$$

where P_v is the projection operator onto \mathcal{H}_v . This means that almost all of $|\psi\rangle$ lies in \mathcal{H}_v . (In CSM, a phase point is always entirely within some macro-state.)

- (7) Thermal equilibrium: Typically, there is a dominant macro-space \mathcal{H}_{eq} that has a dimension that is almost equal to D :

$$\frac{\dim \mathcal{H}_{eq}}{\dim \mathcal{H}} \approx 1. \quad (10)$$

A system with wave function ψ is in equilibrium if the wave function ψ is very close to \mathcal{H}_{eq} in the sense of Equation (9): $\langle \psi | P_{eq} | \psi \rangle \approx 1$. Here’s a simple example: Consider a gas consisting of $n = 10^{23}$ atoms in a box $\Lambda \subseteq \mathbb{R}^3$. The system is governed by quantum mechanics. We orthogonally decompose the Hilbert space \mathcal{H} into fifty-one macro-spaces: $\mathcal{H}_0 \oplus \mathcal{H}_2 \oplus \mathcal{H}_4 \oplus \dots \oplus \mathcal{H}_{100}$, where \mathcal{H}_v is the subspace corresponding to the macro-state such that the number of atoms in the left half of the box is between $(v - 1)\%$ and $(v + 1)\%$ of n , with the endpoints being the exceptions: \mathcal{H}_0 is the interval 0%–1%, and \mathcal{H}_{100} is the interval 99%–100%. In this example, \mathcal{H}_{50} has the overwhelming majority of dimensions and is thus the equilibrium macro-space. A system whose wave function is very close to \mathcal{H}_{50} is in equilibrium (for this choice of macro-states).

- (8) Boltzmann entropy: The Boltzmann entropy of a quantum-mechanical system with wave function ψ that is very close to a macro-state \mathcal{H}_v is given by

$$S_B(\psi) = k_B \log(\dim \mathcal{H}_v), \quad (11)$$

where \mathcal{H}_v denotes the subspace containing almost all of ψ in the sense of Equation (9). The thermal equilibrium state thus has the maximum entropy:

$$S_B(eq) = k_B \log(\dim \mathcal{H}_{eq}) \approx k_B \log(D), \quad (12)$$

where \mathcal{H}_{eq} denotes the equilibrium macro-state. (In CSM, Boltzmann entropy of a phase point is proportional to the logarithm of the volume measure of the macro-state it belongs to.)

- (9) Low-entropy initial condition: When we consider the universe as a quantum-mechanical system, we postulate a special low-entropy boundary condition on the universal wave function—the quantum-mechanical version of the PH:

$$\Psi(t_0) \in \mathcal{H}_{PH}, \dim \mathcal{H}_{PH} \ll \dim \mathcal{H}_{eq} \approx \dim \mathcal{H}, \quad (13)$$

where \mathcal{H}_{PH} is the PH macro-space with dimension much smaller than that of the equilibrium macro-space.⁷ Hence, the initial state has very low entropy in the sense of Equation (11). (In CSM, the PH says that the initial micro-state is in a low-entropy macro-state with very small volume.)

- (10) A central task of QSM is to establish mathematical results that demonstrate (or suggest) that for μ -most (maybe even all) wave functions, the small subsystems, such as gas in a box, will approach thermal equilibrium.

Above is the individualistic view of QSM in a nutshell. In contrast, the ensembler view of QSM differs in several ways. First, on the ensembler view, instead of focusing on the wave function of an individual system, the focus is on an ensemble of systems that have the same statistical state \hat{W} , a density matrix.⁸

It evolves according to the von Neumann equation:

$$i\hbar \frac{d\hat{W}(t)}{dt} = [\hat{H}, \hat{W}]. \quad (14)$$

The crucial difference between the individualistic and the ensembler views of QSM lies in the definition of thermal equilibrium. On the ensembler view, a system is in thermal equilibrium if:

$$W = \rho_{mc} \text{ or } W = \rho_{can}, \quad (15)$$

where ρ_{mc} is the micro-canonical ensemble and ρ_{can} is the canonical ensemble.⁹

⁷ We should assume that \mathcal{H}_{PH} is finite-dimensional, in which case we can use the normalized surface area measure on the unit sphere as the typicality measure for point (10) in Section 2.2. It remains an open question in QSM about how to formulate the low-entropy initial condition when the initial macro-space is infinite-dimensional.

⁸ Ensemblers would further insist that it makes no sense to talk about the thermodynamic state of an individual system.

⁹ The micro-canonical ensemble is the projection operator onto the energy shell \mathcal{H} normalized by its dimension. The canonical ensemble is:

$$\rho_{can} = \frac{\exp(-\beta \hat{H})}{Z}, \quad (16)$$

where $Z = \text{tr} \exp(-\beta \hat{H})$, and β is the inverse temperature of the quantum system.

For the QSM individualist, if the micro-state ψ of a system is close to some macro-space \mathcal{H}_v in the sense of Equation (9), we can say that the macro-state of the system is \mathcal{H}_v . The macro-state is naturally associated with a density matrix:

$$\hat{W}_v = \frac{I_v}{\dim \mathcal{H}_v}, \quad (17)$$

where I_v is the projection operator onto \mathcal{H}_v . \hat{W}_v is thus a representation of the macro-state. It can be decomposed into wave functions, but the decomposition is not unique. Different measures can give rise to the same density matrix. One such choice is $\mu(d\psi)$, the uniform distribution over wave functions:

$$\hat{W}_v = \int_{S(\mathcal{H}_v)} \mu(d\psi) |\psi\rangle \langle \psi|. \quad (18)$$

In Equation (18), \hat{W}_v is defined with a choice of measure on wave functions in \mathcal{H}_v . However, we should not be misled into thinking that the density matrix is derived from wave functions. What is intrinsic to a density matrix is its geometrical meaning in the Hilbert space. In the case of \hat{W}_v , as shown in the canonical description of Equation (17), it is just a normalized projection operator.¹⁰

3. Density-Matrix Realism

According to wave-function realism, the quantum state of the universe is objective and pure. On this view, Ψ is both the micro-state of QSM and a dynamical object of quantum mechanics.

Let us recall the arguments for wave-function realism. Why do we attribute objective status to the quantum state represented by a wave function? It is because the wave function plays crucial roles in the realist quantum theories. In BM, the wave function appears in the fundamental dynamical equations and guides particle motion. In GRW, the wave function spontaneously collapses and gives rise to macroscopic configurations of tables and chairs. In EQM, the wave function is the whole world. If the universe is accurately described by BM, GRW, or EQM, then the wave function is an active ‘agent’ that makes a difference in the world. The wave function cannot represent just our ignorance. It has to be objective, so the arguments go. But what is the nature of the quantum state that it represents? As mentioned in the beginning of this article, there are several interpretations: the two field interpretations, the nomological interpretation, and the *sui generis* interpretation.

On the other hand, we often use W , a density matrix, to represent our ignorance of ψ , the actual wave function of a quantum system. W can also represent a macro-state in QSM.¹¹

¹⁰ Thanks to Sheldon Goldstein for helping me appreciate the intrinsic meaning of density matrices. That was instrumental in the final formulation of the IPH in Section 4.2.

¹¹ In some cases, W is easier for calculation than Ψ , such as in the case of GRW collapse theories where there are multiple sources of randomness. Thanks to Roderich Tumulka for discussions here.

Is it possible to be a realist about the density matrix of the universe and attribute objective status to the quantum state it represents? That depends on whether we can write down realist quantum theories directly in terms of W . Perhaps W does not have enough information to be the basis of a realist quantum theory. However, if we can formulate quantum dynamics directly in terms of W instead of Ψ such that W guides Bohmian particles, or W collapses, or W realizes the emergent multiverse, then we will have good reasons for taking W to represent something objective in those theories. At the very least, the reasons for that will be on a par with those for wave-function realism in the Ψ -theories.

However, can we describe the quantum universe with W instead of Ψ ? The answer is yes. Dürr *et al.* ([2005]) have worked out the Bohmian version. In this section, I describe how. Let us call this new framework density-matrix realism.¹² I will use W-Bohmian mechanics (W-BM) as the main example and explain how a fundamental density matrix can be empirically adequate for describing a quantum world. We can also construct W-Everett theories and W-Ghirardi–Rimini–Weber (W-GRW) theories. Similar to wave-function realism, density-matrix realism is open to several interpretations. At the end of this section, I will provide three field interpretations of W . In Section 6, I discuss and motivate a nomological interpretation.

3.1. W-Bohmian mechanics

First, we illustrate the differences between wave-function realism and density-matrix realism by thinking about two different Bohmian theories.

In standard BM, an N -particle universe at a time t is described by $(Q(t), \Psi(t))$. The universal wave function guides particle motion and provides the probability distribution of particle configurations. Given the centrality of Ψ in BM, we take the wave function to represent something objective (and it is open to several realist interpretations).

It is somewhat surprising that we can formulate a Bohmian theory with only W and Q . This was introduced as W-BM in (Dürr *et al.* [2005]). The fundamental density matrix $W(t)$ is governed by the von Neumann equation, Equation (14). Next, the particle configuration $Q(t)$ evolves according to an analogue of the guidance equation (W-guidance equation):

$$\frac{dQ_i}{dt} = \frac{\hbar}{m_i} \operatorname{Im} \frac{\nabla_q W(q, q', t)}{W(q, q', t)} (q = q' = Q). \quad (19)$$

¹² The possibility that the universe can be described by a fundamental density matrix (mixed state) has been suggested by multiple authors and explored to various extents (see Footnote 2). What is new in this article is the combination of density-matrix realism with the IPH (Section 4) and the argument for the nomological thesis (Section 6) based on that. However, density-matrix realism is unfamiliar enough to warrant some clarifications and developments.

(Here we have set aside spin degrees of freedom. If we include spin, we can add the partial trace operator tr_c before each occurrence of W .) Finally, we can impose an initial probability distribution similar to that of the quantum equilibrium distribution:

$$P(Q(t_0) \in dq) = W(q, q, t_0) dq. \quad (20)$$

The system is also equivariant: if the probability distribution holds at t_0 , it holds at all times.¹³

With the defining equations—the von Neumann equation, Equation (14), and the W-guidance equation, Equation (19)—and the initial probability distribution, Equation (20), we have a theory that directly uses a density matrix $W(t)$ to characterize the trajectories $Q(t)$ of the universe's N particles. If a universe is accurately described by W-BM, then W represents the fundamental quantum state in the theory that guides particle motion; it does not do so via some other entity Ψ . If we have good reasons to be a wave-function realist in BM, then we have equally good reasons to be a density-matrix realist in W-BM.

W-BM is empirically equivalent to BM with respect to the observable quantum phenomena, that is, pointer readings in quantum-mechanical experiments. By the usual typicality analysis (Dürr *et al.* [1992]), this follows from Equation (20), which is analogous to the quantum equilibrium distribution in BM. With the respective dynamical equations, both BM and W-BM generate an equivariant Born-rule probability distribution over all measurement outcomes.¹⁴

3.2. W-Everettian and W-Ghirardi–Rimini–Weber theories

W-BM is a quantum theory in which the density matrix is objective. In this theory, realism about the universal density matrix is based on the central role it plays in the laws of a W-Bohmian universe: it appears in the fundamental dynamical equations and it guides particle motion. (In Section 3.3, we will provide three concrete physical interpretations of W .) What about other quantum theories, such as Everettian and GRW theories? Is it possible to ‘replace’ their universal wave functions with universal

¹³ Equivariance holds because of the following continuity equation:

$$\frac{\partial W(q, q, t)}{\partial t} = -\text{div}(W(q, q, t)v),$$

where v denotes the velocity field generated via Equation (19); see (Dürr *et al.* [1992], [2005]).

¹⁴ Here I am assuming that two theories are empirically equivalent if they assign the same probability distribution to all possible outcomes of experiments. This is the criterion used in the standard Bohmian statistical analysis (Dürr *et al.* [1992]). Empirical equivalence between BM and W-BM follows from the equivariance property plus the quantum equilibrium distribution. Suppose W-BM is governed by a universal density matrix W and suppose BM is governed by a universal wave function chosen at random whose statistical density matrix is W . Then the initial particle distributions on both theories are the same: $W(q, q, t_0)$. By equivariance, the particle distributions will always be the same. Hence, they always agree on what is typical (see Dürr *et al.* [2005]). This is a general argument. A more detailed subsystem statistical analysis, in terms of conditional density matrices, would be illuminating. A first step in this direction can be found in (Chen [unpublished-c]).

density matrices? We show that such suggestions are also possible.¹⁵ First, let us define local beables (in the sense of Bell [2004]). Local beables are the part of the ontology that is localized (to some bounded region) in physical space. Neither the total energy function nor the wave function is a local beable. Candidate local beables include particles, spacetime events (flashes), and matter density ($m(x, t)$).

For the Everettian theory with no local beables (S0), we can postulate that the fundamental quantum state is represented by a density matrix $W(t)$ that evolves unitarily by the von Neumann equation, Equation (14). Let us call this theory W-Everett theory (W-S0). Since there are no additional variables in the theory, the density matrix represents the entire quantum universe. The density matrix will give rise to many branches that (for all practical purposes) do not interfere with each other. The difference is that there will be (in some sense) more branches in the W-Everett quantum state than in the Everett quantum state. In the W-Everett universe, the world history will be described by the undulation of the density matrix.¹⁶

It is difficult to find tables and chairs in a universe described only by a quantum state. One proposal is to add ‘local beables’ to the theory in the form of a mass-density ontology $m(x, t)$. The wave-function version was introduced as ‘Sm’ by Allori *et al.* ([2011]). The idea is that the wave function evolves by the Schrödinger equation and determines the shape of the mass density. This idea can be used to construct a density-matrix version (W-Sm). In this theory, $W(t)$ will evolve unitarily by the von Neumann equation. Next, we can define the mass-density function directly in terms of $W(t)$:

$$m(x, t) = \text{tr}(M(x)W(t)), \tag{21}$$

where x is a physical space variable, $M(x) = \sum_i m_i \delta(Q_i - x)$ is the mass-density operator, which is defined via the position operator $Q_i \psi(q_1, q_2, \dots, q_n) = q_i \psi(q_1, q_2, \dots, q_n)$. This allows us to determine the mass-density ontology at time t via $W(t)$.

For the density-matrix version of GRW theory with just a quantum state (W-GRW0), we need to introduce the collapse of a density matrix. Similar to the wave function in GRW0, between collapses, the density matrix in W-GRW0 will evolve unitarily according to the von Neumann equation. It collapses randomly, where the random time for an N -particle system is distributed with rate $N\lambda$, where λ is of order 10^{-15} s^{-1} . At a random time when a collapse occurs at ‘particle’ k , the post-collapse density matrix is the following:

$$W_{T^+} = \frac{\Lambda_k(X)^{1/2} W_T \Lambda_k(X)^{1/2}}{\text{tr}(W_T \Lambda_k(X))}. \tag{22}$$

¹⁵ Thanks to Roderich Tumulka, Sheldon Goldstein, and Matthias Lienert for discussions here. The W-GRW formalism was first suggested by Allori *et al.* ([2013]).

¹⁶ W-S0 is a novel version of Everettian theory, one that will require more mathematical analysis to fully justify the emergence of macroscopic branching structure. It faces the familiar preferred-basis problem as standard Everett does. In addition, on W-S0 there will be some non-uniqueness in the decompositions of the Hilbert space into macro-spaces. I leave the analysis for future work.

Here, X is distributed by the following probability density:

$$\rho(x) = \text{tr}(W_{T^-} \Lambda_i(x)), \tag{23}$$

where W_{T^-} is the post-collapse density matrix, W_{T^+} is the pre-collapse density matrix, X is the centre of the actual collapse, and $\Lambda_i(x)$ is the collapse rate operator.¹⁷

For the GRW theory (W-GRWm) with both a quantum state $W(t)$ and a mass-density ontology $m(x, t)$, we can combine the above steps: $W(t)$ evolves by the von Neumann equation that is randomly interrupted by collapses (Equation (22)) and $m(x, t)$ is defined by Equation (21). We can define GRW with a flash-ontology (W-GRWf) in a similar way, by using $W(t)$ to characterize the distribution of flashes in physical spacetime. The flashes are the spacetime events at the centres (X) of the W-GRW collapses.

To sum up, in W-S0, the entire world history is described by $W(t)$; in W-Sm, the local beables (mass-density) is determined by $W(t)$; in W-GRW theories, $W(t)$ spontaneously collapses. These roles were originally played by Ψ , and now they are played by W . In so far as we have good reasons for wave-function realism based on the roles that Ψ plays in the Ψ -theories, we have equally good reasons for density-matrix realism if the universe is accurately described by W-theories.

3.3. Field interpretations of W

Realism about the density matrix only implies that it is objective and not epistemic. Realism is compatible with a wide range of concrete interpretations of what the density matrix represents. In this section, I provide three field interpretations of the density matrix. But they do not exhaust all available options. In Section 6, I motivate a nomological interpretation of the density matrix that is also realist.

In debates about the metaphysics of the wave function, realists have offered several interpretations of Ψ . Wave-function realists, such as Albert and Loewer, have offered a concrete physical interpretation: Ψ represents a physical field on the high-dimensional configuration space that is taken to be the fundamental physical space.¹⁸

¹⁷ A collapse rate operator is defined as follows:

$$\Lambda_i(x) = \frac{1}{(2\pi\sigma^2)^{3/2}} e^{-\frac{Q_k - x^2}{2\sigma^2}},$$

where Q_k is the position operator of ‘particle’ k , and σ is a new constant of nature of order 10^{-7} m postulated in current GRW theories. Compare W-GRW to Ψ -GRW, where collapses happen at the same rate, and the post-collapse wave function is the following:

$$\Psi_{T^+} = \frac{\Lambda_i(X)^{1/2} \Psi_{T^-}}{\|\Lambda_i(X)^{1/2} \Psi_{T^-}\|}, \tag{24}$$

with the collapse centre X being chosen randomly with probability distribution $\rho(x) = \|\Lambda_i(x)^{1/2} \Psi_{T^-}\|^2 dx$.

¹⁸ In (Chen [2017]), I argue against this view and suggest that there are many good reasons—internal and external to quantum mechanics—for taking the low-dimensional physical spacetime to be fundamental.

Can we interpret the density matrix in a similar way? Let us start with a mathematical representation of the density matrix $W(t)$. It is defined as a positive, bounded, self-adjoint operator $\hat{W} : \mathcal{H} \rightarrow \mathcal{H}$ with $\text{tr} \hat{W} = 1$. For W-BM, the configuration space \mathbb{R}^{3N} , and a density operator \hat{W} , the relevant Hilbert space is \mathcal{H} , which is a subspace of the total Hilbert space, that is, $\mathcal{H} \subseteq \mathcal{H}_{total} = L^2(\mathbb{R}^{3N}, \mathbb{C})$. Now, the density matrix \hat{W} can also be represented as a function

$$W : \mathbb{R}^{3N} \times \mathbb{R}^{3N} \rightarrow \mathbb{C}. \quad (25)$$

(If we include spin, the range will be the endomorphism space $\text{End}(\mathbb{C}^k)$ of the space of linear maps from \mathbb{C}^k to itself. Notice that we have already used the position representation in Equation (19) and Equation (20).)

This representation enables three field interpretations of the density matrix. Let us use W-BM as an example. First, the fundamental space is represented by \mathbb{R}^{6N} , and W represents a field on that space that assigns properties (represented by complex numbers) to each point in \mathbb{R}^{6N} . In the Bohmian version, W guides the motion of a ‘world particle’ like a river guides the motion of a ping pong ball. (However, the world particle only moves in a \mathbb{R}^{3N} subspace.) Second, the fundamental space is \mathbb{R}^{3N} , and W represents a multi-field on that space that assigns properties to every ordered pair of points (q, q') in \mathbb{R}^{3N} . The world particle moves according to the gradient taken with respect to the first variable of the multi-field. Third, the fundamental space is the physical space represented by \mathbb{R}^3 , and the density matrix represents a multi-field that assigns properties to every ordered pair of N -regions, where each N -region is composed of N points in physical space. On this view, the density matrix guides the motion of N particles in physical space.¹⁹

These three field interpretations are available to the density-matrix realists. In so far as we have good grounds for accepting the field interpretations of wave-function realism, we have equally good grounds for accepting these interpretations for the W-theories. These physical interpretations, I hope, can provide further reasons for wave-function realists to take seriously the idea that density matrices can represent something physically significant. In Section 6, we introduce a new interpretation of W as something nomological, and we will motivate that with the new IPH. That, I believe, is the most interesting realist interpretation of the universal density matrix all things considered.

4. The Initial Projection Hypothesis

W-quantum theories are alternatives to Ψ -quantum theories. However, all of these theories are time-symmetric, as they obey time-reversal invariance.

¹⁹ For discussions about the multi-field interpretation, see (Forrest [1988]; Belot [2012]; Chen [2017], [unpublished-b], Section 3; Hubert and Romano [2018]).

In statistical mechanics, a fundamental postulate is added to the time-symmetric dynamics: the PH, which is a low-entropy boundary condition of the universe. In this section, we will first discuss the wave-function version of the PH. Then we will use it to pick out a special density matrix. I call this the ‘initial projection hypothesis’. Finally, we point out some connections between the IPH and Penrose’s Weyl curvature hypothesis.

4.1. The past hypothesis

The history of the PH goes back to Boltzmann.²⁰ To explain time-asymmetry in a universe governed by time-symmetric equations, Boltzmann’s solution is to add a boundary condition: the universe started in a special state of very low-entropy. Feynman agrees: ‘For some reason, the universe at one time had a very low entropy for its energy content, and since then the entropy has increased’.²¹ Such a low-entropy initial condition explains the arrow of time in thermodynamics.²²

Albert ([2000]) has called this condition the ‘past hypothesis’. However, his proposal is stronger than the usual one concerning a low-entropy initial condition. The usual one just postulates that the universe started in some low-entropy macro-state. It can be any of the many macro-states, so long as it has sufficiently low entropy. Albert’s PH postulates that there is a particular low-entropy macro-state that the universe starts in—the one that underlies the reliability of our inferences to the past. It is the task of cosmology to discover that initial macro-state. In what follows, I refer to the strong version of PH unless indicated otherwise.²³

In QSM, PH takes the form of point (9) in Section 2.2. That is, the micro-state (a wave function) starts in a particular low-dimensional subspace in Hilbert space (the PH-subspace). However, it does not pin down a unique micro-state. There is still a continuous infinity of possible micro-states compatible with the PH-subspace.

It is plausible to think that, for PH to work as a successful explanation for the second law, it has to be on a par with other fundamental laws of nature. That is, we should take PH to be a law of nature and not just a contingent initial condition, for otherwise it might be highly unlikely that our past was in lower entropy and that our inferences to the past are reliable. Already in the context of a weaker version of PH, Feynman ([2017]) suggests that the low-entropy initial condition should be understood as a law of nature. However, PH by itself is not enough. Since there are anti-thermodynamic

²⁰ For an extended discussion, see (Albert [2000]; Callender [2011]; Boltzmann [2012]).

²¹ See (Feynman *et al.* [2015], pp. 46–8).

²² See (Penrose [1979]; Ehrenfest and Ehrenfest [2002]; Lebowitz [2008]) for more discussions about a low-entropy initial condition. See (Earman [2006]) for worries about the PH as an initial condition for the universe. See (Goldstein *et al.* [2016]) for a discussion about the possibility, and some recent examples, of explaining the arrow of time without the PH.

²³ In (Chen [forthcoming]), a companion paper, I discuss different versions of the PH—the strong, the weak, and the fuzzy—as well as their implications for the uniqueness of the initial quantum state that we will come to soon. The upshot is that in all cases it will be sufficiently unique for eliminating statistical-mechanical probabilities.

exceptions even for trajectories starting from the PH-subspace, it is crucial to impose another law about a uniform probability distribution on the subspace. This is the quantum analogue of what Albert ([2000]) calls the statistical postulate (SP). It corresponds to the normalized surface area measure we specified in point (4) in Section 2.2. We used it to state the typicality statement in point (10). Loewer calls the joint system—the package of laws that includes PH and SP in addition to the dynamical laws of physics—the Mentaculus vision.²⁴

4.2. Introducing the initial projection hypothesis

The PH uses a low-entropy macro-state (PH-subspace) to constrain the micro-state of the system (a state vector in QSM). This is natural from the perspective of wave-function realism, according to which the state vector (the wave function) represents the physical degrees of freedom of the system. The initial state of the system is described by a normalized wave function $\Psi(t_0)$. $\Psi(t_0)$ has to lie in the special low-dimensional Hilbert space \mathcal{H}_{PH} with $\dim\mathcal{H}_{PH} \ll \dim\mathcal{H}_{eq}$. Moreover, there are many different choices of initial wave functions in \mathcal{H}_{PH} . That is, PH is compatible with many different low-entropy wave functions. Furthermore, for stating the typicality statements, we also need to specify a measure μ on the unit sphere of \mathcal{H}_{PH} . For the finite-dimensional case, it is just the normalized surface area measure.

Density-matrix realism suggests an alternative way to think about the low-entropy boundary condition. PH pins down the initial macro-state \mathcal{H}_{PH} , a special subspace of the total Hilbert space. Although \mathcal{H}_{PH} is compatible with many density matrices, there is a natural choice—the normalized projection operator onto \mathcal{H}_{PH} . Just as in Equation (17), we can specify it as:

$$\hat{W}_{IPH}(t_0) = \frac{I_{PH}}{\dim\mathcal{H}_{PH}}, \quad (26)$$

where t_0 represents a temporal boundary of the universe, I_{PH} is the projection operator onto \mathcal{H}_{PH} , \dim counts the dimension of the Hilbert space, and $\dim\mathcal{H}_{PH} \ll \dim\mathcal{H}_{eq}$. Since the quantum state at t_0 has the lowest entropy, we call t_0 the initial time. We shall call Equation (26) the initial projection hypothesis (IPH). In words: the initial density matrix of the universe is the normalized projection onto the PH-subspace.

I propose that we add IPH to any W-quantum theory. The resultant theories will be called W_{IPH} -theories. For example, here are the equations of W_{IPH} -BM:

²⁴ For developments and defences of the nomological account of the PH and the SP, see (Albert [2000]; Loewer [2007]; Wallace [unpublished], [2012]; Loewer [unpublished]). Albert and Loewer are writing mainly in the context of CSM. The Mentaculus vision is supposed to provide a ‘probability map of the world’. As such, it requires one to take the probability distribution very seriously.

To be sure, the view that PH is nomological and explanatorily fundamental has been debated; see, for example, (Price [1996], [2004]; Callender [2004]) for discussion. However, these arguments are no more threatening to IPH being a law than PH being a law. We will come back to this point after introducing IPH.

- (i) $\hat{W}_{IPH}(t_0) = \frac{I_{PH}}{\dim \mathcal{H}_{PH}}$,
(ii) $P(Q(t_0) \in dq) = W_{IPH}(q, q, t_0) dq$,
(iii) $i\hbar \frac{\partial \hat{W}}{\partial t} = [\hat{H}, \hat{W}]$,
(iv) $\frac{dQ_i}{dt} = \frac{\hbar}{m_i} \text{Im} \frac{\nabla_{q_i} W_{IPH}(q, q', t)}{W_{IPH}(q, q', t)} (q = q' = Q)$.

(i) is IPH and (ii)–(iv) are the defining equations of W-BM. (Given the initial quantum state $\hat{W}_{IPH}(t_0)$, there is a live possibility that for every particle at t_0 , its velocity is zero. However, even in this possibility, as long as the initial quantum state ‘spreads out’ later, as we assume it would, the particle configuration will typically start moving at a later time. This is true because of equivariance.²⁵)

Contrast these equations with BM formulated with wave functions and PH (not including SP for now), which will be called Ψ_{PH} -BM:

- (i') $\Psi(t_0) \in \mathcal{H}_{PH}$,
(ii') $P(Q(t_0) \in dq) = |\Psi(q, t_0)|^2 dq$,
(iii') $i\hbar \frac{\partial \Psi}{\partial t} = \hat{H}\Psi$,
(iv') $\frac{dQ_i}{dt} = \frac{\hbar}{m_i} \text{Im} \frac{\nabla_{q_i} \Psi(q, t)}{\Psi(q, t)} (Q)$.

IPH (i) in W_{IPH} -BM plays the same role as PH (i') in Ψ_{PH} -BM. Should IPH be interpreted as a law of nature in W_{IPH} -theories? I think it should be, for the same reason that PH should be interpreted as a law of nature in the corresponding theories. The reason that PH should be interpreted as a law is because it is a particularly simple and informative statement that accounts for the widespread thermodynamic asymmetry in time.²⁶ PH is simple because it characterizes a simple macro-state \mathcal{H}_{PH} , of which the initial wave function is a vector. PH is informative because with PH the dynamical equations predict time-asymmetry and without PH the dynamical equations cannot. Similarly, IPH is simple because it provides crucial resources for explaining the arrow of time. IPH is informative because it is essential for explaining the time-asymmetry in a quantum universe described by a density matrix. (This is in addition to the fact that IPH helps determine the W_{IPH} -version of the guidance equation, (iv).) To be sure, PH and IPH as laws face the same worries: both are statements about boundary conditions but we usually think of laws as dynamical equations. However, these worries are no more threatening to IPH being a law than PH being a law.

Let us make three remarks about IPH. First, IPH defines a unique initial quantum state. The quantum state $\hat{W}_{IPH}(t_0)$ is informationally equivalent to the constraint that PH imposes on the initial micro-states. Assuming that PH selects a unique low-entropy macro-state, $\hat{W}_{IPH}(t_0)$ is singled out by the data in PH.²⁷

²⁵ Thanks to Sheldon Goldstein and Tim Maudlin for discussions here.

²⁶ See, for example, (Albert [2000]; Feynman [2017]; Loewer [2007], [unpublished]).

²⁷ The weaker versions of PH are vague about the exact initial low-entropy macro-state. It is vague because even with a choice of macro-variables, there may be many subspaces that can play the role of a low-entropy initial condition. It would be arbitrary, from the viewpoint of wave-function theories, to pick a specific subspace. In contrast, it would not be arbitrary from the viewpoint of W_{PH} -theories, as the specific subspace defines W_{PH} , which determines the dynamics.

Second, on the universal scale, we do not need to impose an additional probability or typicality measure on the Hilbert space. $\hat{W}_{IPH}(t_0)$ is mathematically equivalent to an integral over projection onto each normalized state vectors (wave functions) compatible with PH with respect to a Lebesgue measure. But here we are not defining $\hat{W}_{IPH}(t_0)$ in terms of state vectors. Rather, we are thinking of $\hat{W}_{IPH}(t_0)$ as a geometric object in the Hilbert space: the (normalized) projection operator onto \mathcal{H}_{PH} . That is the intrinsic understanding of the density matrix.²⁸

Third, $\hat{W}_{IPH}(t_0)$ is simple. Related to the first remark, IPH defines $\hat{W}_{IPH}(t_0)$ explicitly as the normalized projection operator onto \mathcal{H}_{PH} . There is a natural correspondence between a subspace and its projection operator. If we specify the subspace, we know what its projection operator is, and vice versa. Since the projection operator onto a subspace carries no more information than that subspace itself, the projection operator is no more complex than \mathcal{H}_{PH} . This is different from Ψ_{PH} , which is confined by PH to be a vector inside \mathcal{H}_{PH} . A vector carries more information than the subspace it belongs to, as specifying a subspace is not sufficient to determine a vector. For example, to determine a vector in an 18-dimensional subspace of a 36-dimensional vector space, we need 18 coordinates in addition to specifying the subspace. The higher the dimension of the subspace, the more information is needed to specify the vector. If PH had fixed Ψ_{PH} (the QSM micro-state), it would have required much more information and become a much more complex posit. PH as it is determines Ψ_{PH} only up to an equivalence class (the QSM macro-state). As we shall see in Section 6, the simplicity of $\hat{W}_{IPH}(t_0)$ will be a crucial ingredient for a new version of the nomological interpretation of the quantum state.

4.3. Connections to the Weyl curvature hypothesis

Let us point out some connections between our IPH and the Weyl curvature hypothesis (WCH) proposed by Penrose ([1979]). Thinking about the origin of the second law of thermodynamics in the early universe with high homogeneity and isotropy, and the relationship between spacetime geometry and entropy, Penrose ([1979], p. 630) proposes a low-entropy hypothesis:

I propose, then, that there should be complete lack of chaos in the initial *geometry*. We need, in any case, some kind of low-entropy constraint on the initial

²⁸ After writing this article, I discovered that Wallace ([forthcoming]) has come to a similar idea in a forthcoming paper. There are some subtle differences. He proposes that we can reinterpret probability distributions in QSM as actual mixed states. Consequently, the problem of statistical-mechanical probability is 'radically transformed' (if not eliminated) in QSM. Wallace's proposal is compatible with different probability distributions and hence different mixed states of the system. It does not require one to choose a particular quantum state such as in Equation (26). Moreover, it is compatible with there being an underlying pure state. In contrast, I propose a particular, natural initial quantum state of the universe based on the PH-subspace—the normalized projection onto the PH-subspace, Equation (26), and there is no underlying pure state. As we discuss in Section 5.1, this also leads to the elimination of statistical-mechanical probability, since the initial state is fixed in the theory. Moreover, as we discuss below, the natural state inherits the simplicity of the PH-subspace, which has implications for the nature of the quantum state. For a more detailed comparison, see (Chen [forthcoming]).

state. But thermal equilibrium apparently held (at least very closely so) for the *matter* (including radiation) in the early stages. So the ‘lowness’ of the initial entropy was not a result of some special matter distribution, but, instead, of some very special initial spacetime geometry. The indications of [previous sections], in particular, are that this restriction on the early geometry should be something like: *the Weyl curvature C_{abcd} vanishes at any initial singularity.*

The Weyl curvature tensor C_{abcd} is the traceless part of the Riemann curvature tensor R_{abcd} . It is not fixed completely by the stress-energy tensor and thus has independent degrees of freedom in Einstein’s general theory of relativity. Since the entropy of the matter distribution is quite high, the origin of thermodynamic asymmetry should be due to the low entropy in geometry, which corresponds very roughly to the vanishing of the Weyl curvature tensor.

WCH is an elegant and simple way of encoding the initial low-entropy boundary condition in the classical spacetime geometry. If WCH could be extended to a quantum theory of gravity, presumably it would pick out a simple subspace (or subspaces) of the total Hilbert space that corresponds to $C_{abcd} \rightarrow 0$. Applying IPH to such a theory, the initial density matrix will be the normalized projection onto that subspace (subspaces).²⁹

5. Theoretical Pay-Offs

W_{IPH} -quantum theories, the result of applying IPH to W-theories, have two theoretical pay-offs, which we explore in this section. These are by no means decisive arguments in favour of the density-matrix framework, but they display some interesting differences with the wave-function framework.

5.1. Harmony between statistical mechanics and quantum mechanics

In W_{IPH} -quantum theories, statistical mechanics is made more harmonious with quantum mechanics. As we pointed out earlier, standard quantum mechanics and QSM contain the wave function in addition to the density matrix, and they require the addition of both the PH and the SP to the dynamical laws. In particular, we have two kinds of probabilities: the quantum-mechanical ones (Born rule probabilities) and the statistical-mechanical ones (SP). The situation is quite different in our framework. This is true for all the W_{IPH} -theories. We will use W_{IPH} -BM ((i)–(iv)) as an example.

W_{IPH} -BM completely specifies the initial quantum state, unlike Ψ_{PH} -BM. For Ψ_{PH} -BM, because of time-reversal invariance, some initial wave functions compatible

²⁹ There is another connection between the current project and Penrose’s work. The W-Everettian theory that we considered in Section 3.2 combined with the IPH is a theory that satisfies strong determinism (Penrose [1999]). This is because the entire history of the W_{IPH} -Everettian universe described by $W_{IPH}(t)$, including its initial condition, is fixed by the laws.

with PH will evolve to lower entropy. These are called ‘anti-entropic’ exceptions. However, the uniform probability distribution (SP) assigns low probability to these exceptions. Hence, we expect that with overwhelming probability the actual wave function is entropic. For W_{IPH} -BM, in contrast, there is no need for something like SP, as there is only one initial density matrix compatible with IPH— $W_{IPH}(t_0)$. It is guaranteed to evolve to future states that have entropic behaviours. Therefore, on the universal scale, W_{IPH} -BM eliminates the need for SP and thus the need for a probability or typicality measure that is in addition to the quantum-mechanical measure (ii). This is a nice feature of W_{IPH} -theories, as it is desirable to unify the two sources of randomness: quantum mechanical and statistical mechanical. Of course, wave functions and statistical-mechanical probabilities are still useful to analyse subsystems such as gas in a box, but they no longer play fundamental roles in W_{IPH} -theories. Another strategy to eliminate SP has been explored in the context of GRW jumps by Albert ([2000]). Wallace ([unpublished], [2012]) has proposed a replacement of SP with a non-probabilistic constraint on the micro-state, giving rise to the simple dynamical conjecture. These are quite different proposals, all of which deserve further developments.

5.2. Descriptions of the universe and the subsystems

W_{IPH} -quantum theories also bring more unity to the kinematics and the dynamics of the universe and the subsystems.

Let us start with a quantum-mechanical universe U . Suppose it contains many subsystems. Some of them will be interacting heavily with the environment, while others will be effectively isolated from the environment. For a universe that contains some quasi-isolated subsystems (interactions with the environment effectively vanish), the following is a desirable property:

Dynamic Unity: The dynamical laws of the universe are the same as the effective laws of most quasi-isolated subsystems.

Dynamic unity is a property that can come in degrees, rather than an ‘on-or-off’ property. Theory A has more dynamic unity than Theory B, if the fundamental equations in A are valid in more subsystems than those in B. This property is desirable, but not indispensable. It is desirable because law systems that apply both at the universal level and at the subsystem level are unifying and explanatory.

W -BM has more dynamic unity than BM formulated with a universal wave function. For quantum systems without spin, we can always follow Dürr *et al.* ([1992]) to define conditional wave functions in BM. For example, if the universe is partitioned into a system S_1 and its environment S_2 , then for S_1 , we can define its conditional wave function:

$$\psi_{cond}(q_1) = C\Psi(q_1, Q_2), \quad (27)$$

where C is a normalization factor and Q_2 is the actual configuration of S_2 . $\psi_{cond}(q_1)$ always gives the velocity field for the particles in S_1 according to the guidance equation. However, for quantum systems with spin, this is not always true. Since BM is described by $(\Psi(t), Q(t))$, it does not contain actual values of spin. Since there are no actual spins to plug into the spin indices of the wave function, we cannot always define conditional wave functions in an analogous way. Nevertheless, in those circumstances, we can follow Dürr *et al.* ([2005]) to define a conditional density matrix for S_1 , by plugging in the actual configuration of S_2 and tracing over the spin components in the wave function associated with S_2 .³⁰ The conditional density matrix will guide the particles in S_1 by the W-guidance equation (the spin version with the partial trace operator).

In W-BM, the W-guidance equation is always valid for the universe and the subsystems. In BM, sometimes subsystems do not have conditional wave functions, and thus the wave-function version of the guidance equation is not always valid. In this sense, the W-BM equations are valid in more circumstances than the BM equations. However, this point does not rely on IPH.

What about Everettian and GRW theories? Since GRW and Everettian theories do not have fundamental particles, we cannot obtain conditional wave functions for subsystems as in BM. However, even in the Ψ -versions of GRW and Everett, many subsystems will not have pure-state descriptions by wave functions due to the prevalence of entanglement. Most subsystems can be described only by a mixed-state density matrix, even when the universe as a whole is described by a wave function. In contrast, in W_{IPH} -Everett theories and W_{IPH} -GRW theories, there is more uniformity across the subsystem level and the universal level: the universe as a whole as well as most subsystems are described by the same kind of object—a (mixed-state) density matrix. Since state descriptions concern the kinematics of a theory, we say that W-Everett and W-GRW theories have more kinematic unity than their Ψ -counterparts:

Kinematic Unity: The state description of the universe is of the same kind as the state descriptions of most quasi-isolated subsystems.

So far, my main goal has been to show that density-matrix realism + IPH is a viable position. They have theoretical pay-offs that are interestingly different from those in the original package (wave-function realism + PH). In the next section, we look at their relevance to the nature of the quantum state.

³⁰ The conditional density matrix for S_1 is defined as:

$$W_{cond S_1' S_1}(q_1, q_1') = \frac{1}{N} \sum_{s_2} \Psi^{s_1 s_2}(q_1, Q_2) \Psi_{s_1 s_2}^*(q_1', Q_2). \tag{28}$$

Here, the normalizing factor is:

$$N = \int_{Q_1} dq_1 \sum_{s_1 s_2} \Psi^{s_1 s_2}(q_1, Q_2) \Psi_{s_1 s_2}^*(q_1', Q_2). \tag{29}$$

6. The Nomological Thesis

Combining density-matrix realism with IPH gives us W_{IPH} -quantum theories that have interesting theoretical pay-offs. We have also argued that the initial quantum state in such theories would be simple and unique. In this section, we show that the latter fact lends support to the nomological interpretation of the quantum state:

The Nomological Thesis: The initial quantum state of the world is nomological.

However, ‘nomological’ has several senses and has been used in several ways in the literature. We will start with some clarifications.

6.1. The classical case

We can clarify the sense of the ‘nomological’ by taking another look at classical mechanics. In classical N -particle Hamiltonian mechanics, it is widely accepted that the Hamiltonian function is nomological, and that the ontology consists in particles with positions and momenta. Their state is given by $X = (q_1(t), \dots, q_N(t); p_1(t), \dots, p_n(t))$, and the Hamiltonian is $H = H(X)$. Particles move according to the Hamiltonian equations:

$$\frac{dq_i(t)}{dt} = \frac{\partial H}{\partial p_i}, \quad \frac{dp_i(t)}{dt} = -\frac{\partial H}{\partial q_i}. \quad (30)$$

Their motion corresponds to a trajectory in phase space. The velocity field on phase space is obtained by taking suitable derivatives of the Hamiltonian function H . The equations have the form:

$$\frac{dX}{dt} = F(X) = F^H(X). \quad (31)$$

Here, $F^H(X)$ is $H(q, p)$ with suitable derivative operators. The Hamiltonian equations have a simple form, because H is simple. H can be written explicitly as follows:

$$H = \sum_i^N \frac{p_i^2}{2m_i} + V. \quad (32)$$

V takes on this form when we consider electric and gravitational potentials:

$$V = \frac{1}{4\pi\epsilon_0} \sum_{1 \leq j \leq k \leq N} \frac{e_j e_k}{|q_j - q_k|} + \sum_{1 \leq j \leq k \leq N} \frac{Gm_j m_k}{|q_j - q_k|}. \quad (33)$$

That is, the right-hand side of the Hamiltonian equations, after making the Hamiltonian function explicit, are still simple. H is just a convenient shorthand for Equation (32) and Equation (33). Moreover, H is also fixed by the theory. A classical universe is governed by the dynamical laws plus the fundamental interactions. If H were different in Equation (31), then we would have a different physical theory

(though it would still belong to the class of theories called classical mechanics). For example, we can add another term in Equation (33) to encode another fundamental interaction, which will result in a different theory.

Consequently, it is standard to interpret H as a function in Equation (30) that does not represent things or properties of the ontology. Expressed in terms of H , the equations of motion take a particularly simple form. The sense that H is nomological is that (i) it generates motion, (ii) it is simple, (iii) it is fixed by the theory (nomologically necessary), and (iv) it does not represent things in the ontology. In contrast, the position and momentum variables in Equation (30) are ‘ontological’ in that they represent things and properties of the ontology, take on complicated values, change according to H , and are not completely fixed by the theory (contingent).

6.2. The quantum case

It is according to the above sense that Goldstein and Zanghi ([2013]) proposes that the universal wave function in BM is nomological (and governs things in the ontology). With the guidance equation, Ψ generates the motion of particles. It is of the same form as above:

$$\frac{dX}{dt} = F(X) = F^\Psi(X). \quad (34)$$

Why is it simple? Generic wave functions are not simple. However, they observe that, in some formulations of quantum gravity, the universal wave function satisfies the Wheeler–DeWitt equation and is therefore stationary. To be stationary, the wave function does not have time-dependence and probably has many symmetries, in which case it could be quite simple. The Bohmian theory then will explicitly stipulate what the universal wave function is. Therefore, in these theories, provided that Ψ is sufficiently simple, we can afford the same interpretation of Ψ as we can for H in classical mechanics: both are nomological in the above sense.

W_{IPH} -BM also supports the nomological interpretation of the quantum state but via a different route. With the W -guidance equation, W_{IPH} generates the motion of particles. It is of the same form as above:

$$\frac{dX}{dt} = F(X) = F^{W_{IPH}}(X). \quad (35)$$

Why is it simple? Here we do not need to appeal to specific versions of quantum gravity, which are still to be worked out and may not guarantee the simplicity of Ψ . Instead, we can just appeal to IPH. We have argued in Section 4.2 that IPH is simple and that $W_{IPH}(t_0)$ is simple. Since the quantum state evolves unitarily by the von Neumann equation, we can obtain the quantum state at any later time as:

$$\hat{W}_{IPH}(t) = e^{-i\hat{H}t/\hbar} \hat{W}_{IPH}(t_0) e^{i\hat{H}t/\hbar}. \quad (36)$$

Since $W_{IPH}(t)$ is a simple function of the time-evolution operator and the initial density matrix, and since both are simple, $W_{IPH}(t)$ is also simple. So we can think of $W_{IPH}(t)$ just as a convenient shorthand for Equation (36). (This is not true for $|\Psi(t)\rangle = \hat{H} |\Psi(t_0)\rangle$, as generic $|\Psi(t_0)\rangle$ is not simple at all.)

The ‘shorthand’ way of thinking about $W_{IPH}(t)$ implies that the equation of particle motion has a time-dependent form $F^{W_{IPH}}(X, t)$. Does time-dependence undercut the nomological interpretation? It does not in this case, as the $F^{W_{IPH}}(X, t)$ is still simple even with time-dependence. It is true that time-independence is often a hallmark of a nomological object, but it is not always the case. In this case, we have simplicity without time-independence. Moreover, unlike Goldstein and Zanghì ([2013]), we do not need time-independence to argue for the simplicity of the quantum state.

Since $W_{IPH}(t_0)$ is fixed by IPH, $F^{W_{IPH}}$ is also fixed by the theory. Let us expand Equation (35) to make it more explicit:

$$\begin{aligned} \frac{dQ_i}{dt} &= \frac{\hbar}{m_i} \text{Im} \frac{\nabla_q W_{IPH}(q, q', t)}{W_{IPH}(q, q', t)}(Q) \\ &= \frac{\hbar}{m_i} \text{Im} \frac{\nabla_q \langle q | e^{-i\hat{H}t/\hbar} \hat{W}_{IPH}(t_0) e^{i\hat{H}t/\hbar} | q' \rangle}{\langle q | e^{-i\hat{H}t/\hbar} \hat{W}_{IPH}(t_0) e^{i\hat{H}t/\hbar} | q' \rangle} (q = q' = Q). \end{aligned} \tag{37}$$

The initial quantum state (multiplied by the time-evolution operators) generates motion, has a simple form, and is fixed by the boundary condition (IPH) in W_{IPH} -BM. Therefore, it is nomological. This is of course a modal thesis. The initial quantum state, which is completely specified by IPH, could not have been different.

Let us consider other W_{IPH} -theories with local beables. In W_{IPH} -Sm, the initial quantum state has the same simple form and is fixed by IPH. It does not generate a velocity field, since there are no fundamental particles in the theory. Instead, it determines the configuration of the mass-density field on physical space. This is arguably different from the sense of nomological that H in classical mechanics displays. Nevertheless, the mass-density field and the Bohmian particles play a similar role—they are ‘local beables’ that make up tables and chairs, and they are governed by the quantum state. In W_{IPH} -GRWm and W_{IPH} -GRWf, the initial quantum state has the same simple form and is fixed by IPH. It does not generate a velocity field, and it evolves stochastically. This will determine a probability distribution over configurations of local beables—mass densities or flashes—on physical space. The initial quantum state in these theories can be given an extended nomological interpretation, in the sense that condition (i) is extended such that it covers other kinds of ontologies and dynamics: (i') the quantum state determines (deterministically or stochastically) the configuration of local beables.

The W_{IPH} -theories with local beables support the nomological interpretation of the initial quantum state. It can be interpreted in non-Humean ways and Humean ways. On the non-Humean proposal, we can think of the initial quantum state as an additional nomological entity that explains the distribution of particles, fields,

or flashes. On the Humean proposal, in contrast, we can think of the initial quantum state as something that summarizes a separable mosaic. This leads to reconciliation between HS and quantum entanglement.

6.3. Humean supervenience

Recall that according to HS, the 'vast mosaic of local matters of particular fact' is a supervenience base for everything else in the world, the metaphysical ground floor on which everything else depends. On this view, laws of physics are nothing over and above the 'mosaic'. They are just the axioms in the simplest and most informative summaries of the local matters of particular fact. A consequence of HS is that the complete physical state of the universe is determined by the properties and spatiotemporal arrangement of the local matters (suitably extended to account for vector-valued magnitudes) of particular facts. It follows that there should not be any state of the universe that fails to be determined by the properties of individual spacetime points.³¹ Quantum entanglement, if it were in the fundamental ontology, would present an obstacle to HS, because entanglement is not determined by the properties of spacetime points. The consideration above suggests a strong *prima facie* conflict between HS and quantum physics. On the basis of quantum non-separability, Maudlin has proposed an influential argument against HS.³²

W_{IPH} -theories with local beables offer a way out of the conflict between quantum entanglement and HS. A Humean can interpret the laws (including the IPH) as the axioms in the best system that summarize a separable mosaic. Take W_{IPH} -BM as an example:

The W_{IPH} -BM Mosaic: Particle trajectories $Q(t)$ on physical spacetime.

The W_{IPH} -BM Best System: Four equations—the simplest and strongest axioms summarizing the mosaic:

- (i) $\hat{W}_{IPH}(t_0) = \frac{I_{PH}}{\dim T_{t_0}}$,
- (ii) $P(Q(t_0) \in dq) = W_{IPH}(q, q, t_0) dq$,
- (iii) $i\hbar \frac{\partial \hat{W}}{\partial t} = [\hat{H}, \hat{W}]$,
- (iv) $\frac{dQ_i}{dt} = \frac{\hbar}{m_i} \text{Im} \frac{\nabla_q W_{IPH}(q, q', t)}{W_{IPH}(q, q', t)} (q = q' = Q)$.

Notice that (i)–(iv) are simple and informative statements about $Q(t)$. They are expressed in terms of $\hat{W}_{IPH}(t)$, which via law (iii) can be expressed in terms of $\hat{W}_{IPH}(t_0)$. We have argued previously that the initial quantum state can be given a nomological interpretation. The Humean manoeuvre is that the law statements are to be understood as axioms of the best summaries of the mosaic. The mosaic described above is completely separable, while the best system, completely specifying the quantum

³¹ This is one reading of Lewis. Maudlin ([2007]) calls this thesis 'separability'.

³² See (Maudlin [2007], Chapter 2).

state and the dynamical laws, contains all the information about quantum entanglement and superpositions. The entanglement facts are no longer fundamental. As on the original version of HS, the best system consisting of (i)–(iv) supervenes on the mosaic. Hence, this proposal reconciles HS with quantum entanglement. As it turns out, the above version of quantum Humeanism also achieves more theoretical harmony, dynamical unity, and kinematic unity (Section 5), which are desirable from the Humean best-system viewpoint. We can perform similar ‘Humeanization’ manoeuvres on the density matrix in other quantum theories with local beables— W_{IPH} -GRWm, W_{IPH} -GRWf, and W_{IPH} -Sm (although such procedures might not be as compelling).

This version of quantum Humeanism based on W_{IPH} -theories is different from the other approaches in the literature (Albert [1996]; Loewer [1996]; Miller [2014]; Esfeld [2014]; Bhogal and Perry [2015]; Callender [2015]; Esfeld and Decker [2017]). In contrast to the high-dimensional proposal in (Albert [1996]; Loewer [1996]), our version preserves the fundamentality of physical space.

The difference between our version and those in (Miller [2014]; Esfeld [2014]; Bhogal and Perry [2015]; Callender [2015]; Esfeld and Deckert [2017]) is more subtle. They are concerned primarily with Ψ -BM. We can summarize their views as follows (although they do not agree on all the details). There are several parts to their proposals. First, the wave function is merely part of the best system. It is more like parameters in the laws such as mass and charge. Second, just like the rest of the best system, the wave function supervenes on the mosaic of particle trajectories. Third, the wave function does not have to be very simple. The Humean theorizer, on this view, just needs to find the simplest and strongest summary of the particle histories, but the resultant system can be complex simpliciter. One interpretation of this view is that the best system for Ψ_{PH} -BM is just (i')–(iv') in Section 4.2 (although they do not explicitly consider (i')), such that neither the mosaic nor the best system specifies the exact values of the universal wave function. In contrast, our best system completely specifies the universal quantum state. The key difference between our approaches is that their interpretation of the wave function places much weaker constraints than our nomological interpretation does. It is much easier for something to count as being part of the best system on their approach than on ours. While they do not require the quantum state to be simple, we do. For them, the Bohmian guidance equation is likely very complex after plugging in the actual wave function Ψ_{PH} on the right-hand side, but Ψ_{PH} can still be part of their best system.³³ For us, it is crucial that the equation remains simple after plugging in $W_{IPH}(t_0)$ for it to be in the best system. Consequently, $W_{IPH}(t_0)$ is nomological in the sense spelled out in Section 6.1, and we can give it a Humean interpretation similar to that of the Hamiltonian function in CM. Generic Ψ_{PH} , on the other hand, cannot be nomological in our sense. But that is ok for them, as their best-system interpretation does not require the strong nomological condition

³³ See (Dewar [unpublished], Section 5) for some worries about the weaker criterion on the best system.

that we use. Here we do not attempt to provide a detailed comparison; we will do that in future work.

7. Conclusion

I have introduced a new package of views: density-matrix realism, the initial projection hypothesis, and the nomological thesis. In the first two steps, we introduced a new class of quantum theories: W_{IPH} -theories. In the final step, we argue that it is a theory in which the initial quantum state can be given a nomological interpretation. Each is interesting in its own right, and they do not need to be taken together. However, they fit together quite well. They provide alternatives to standard versions of realism about quantum mechanics, a new way to get rid of statistical-mechanical probabilities, and a new solution to the conflict between quantum entanglement and Humean supervenience. To be sure, there are many other features of W_{IPH} -theories in general and the nomological interpretation in particular that are worth exploring further.

The most interesting feature of the new package, I think, is that it brings together the foundations of quantum mechanics and quantum statistical mechanics. In W_{IPH} -theories, the arrow of time becomes intimately connected to the quantum-mechanical phenomena in nature. It is satisfying to see that nature is so unified.

Acknowledgements

I would like to thank the editors and referees of the *British Journal for the Philosophy of Science* for helpful feedback. I am also grateful for stimulating discussions with David Albert, Sheldon Goldstein, and Barry Loewer. I have received helpful feedback from Abhay Ashtekar, Karen Bennett, Harjit Bhogal, Max Bialek, Miren Boehm, Robert Brandenberger, Tad Brennan, Sean Carroll, Juliusz Doboszewski, Detlef Dürr, Denise Dykstra, Michael Esfeld, Veronica Gomez, Hans Halvorson, Harold Hodes, Mario Hubert, Michael Kiessling, Dustin Lazarovici, Stephen Leeds, Matthias Lienert, Niels Linnemann, Chuang Liu, Vera Matarese, Tim Maudlin, Elizabeth Miller, Sebastian Murgueitio, Wayne Myrvold, Jill North, Zee Perry, Davide Romano, Ezra Rubenstein, Charles Sebens, Ted Sider, Joshua Spencer, Noel Swanson, Karim Thébault, Roderich Tumulka, David Wallace, Isaac Wilhelm, Nino Zanghì, and audiences at Cornell University, the University of Western Ontario, the University of Wisconsin-Milwaukee, and the 2018 Rotman Summer Institute in Philosophy of Cosmology.

*Department of Philosophy
University of California, San Diego
San Diego, CA, USA
eddyckm@gmail.com*

References

- Albert, D. Z. [1996]: ‘Elementary Quantum Metaphysics’, in J. T. Cushing, A. Fine and S. Goldstein (eds), *Bohmian Mechanics and Quantum Theory: An Appraisal*, Dordrecht: Kluwer, pp. 277–84.
- Albert, D. Z. [2000]: *Time and Chance*, Cambridge, MA: Harvard University Press.
- Allori, V., Goldstein, S., Tumulka, R. and Zanghì, N. [2011]: ‘Many Worlds and Schrödinger’s First Quantum Theory’, *British Journal for the Philosophy of Science*, **62**, pp. 1–27.
- Allori, V. [2013]: ‘Predictions and Primitive Ontology in Quantum Foundations: A Study of Examples’, *British Journal for the Philosophy of Science*, **65**, pp. 323–52.
- Bell, J. S. [1990]: ‘Against “Measurement”’, *Physics World*, **3**, pp. 33.
- Bell, J. S. [2004]: *Speakable and Unspeakable in Quantum Physics: Collected Papers on Quantum Philosophy*, Cambridge: Cambridge University Press.
- Belot, G. [2012]: ‘Quantum States for Primitive Ontologists’, *European Journal for Philosophy of Science*, **2**, pp. 67–83.
- Bhagal, H. and Perry, Z. [2015]: ‘What the Humean Should Say about Entanglement’, *Noûs*, **51**, pp. 74–94.
- Boltzmann, L. [2012]: *Lectures on Gas Theory*, Chelmsford, MA: Courier Corporation.
- Callender, C. [2004]: ‘Measures, Explanations, and the Past: Should “Special” Initial Conditions Be Explained?’, *British Journal for the Philosophy of Science*, **55**, pp. 195–217.
- Callender, C. [2011]: ‘Thermodynamic Asymmetry in Time’, in E. N. Zalta (ed.), *The Stanford Encyclopedia of Philosophy*, available at <plato.stanford.edu/archives/fall2011/entries/time-thermo/>.
- Callender, C. [2015]: ‘One World, One Beable’, *Synthese*, **192**, pp. 3153–77.
- Chen, E. K. [2017]: ‘Our Fundamental Physical Space: An Essay on the Metaphysics of the Wave Function’, *Journal of Philosophy*, **114**, pp. 333–65.
- Chen, E. K. [2019]: ‘Realism about the Wave Function’ *Philosophy Compass*, **14**, available at <<https://doi.org/10.1111/phc3.12611>>.
- Chen, E. K. [forthcoming]: ‘Time’s Arrow in a Quantum Universe: On the Status of Statistical Mechanical Probabilities’, in V. Allori (ed.), *Statistical Mechanics and Scientific Explanation: Determinism, Indeterminism and Laws of Nature*, Singapore: World Scientific.
- Chen, E. K. [unpublished-a]: ‘The Best Summary of the Quantum World: The Universal Wave Function as a Humean Law’.
- Chen, E. K. [unpublished-b]: ‘An Intrinsic Theory of Quantum Mechanics: Progress in Field’s Nominalistic Program, Part I’, available at <philsci-archive.pitt.edu/14201/>.
- Chen [unpublished-c]: ‘Quantum States of a Time-asymmetric Universe: Wave Function, Density Matrix, and Empirical Equivalence’, available at <arxiv.org/abs/1901.080537>.
- Dewar, N. [unpublished]: ‘La Bohème’, available at <philsci-archive.pitt.edu/13235/>.
- Dürr, D., Goldstein, S., Tumulka, R. and Zanghì, N. [2005]: ‘On the Role of Density Matrices in Bohmian Mechanics’, *Foundations of Physics*, **35**, pp. 449–67.
- Dürr, D., Goldstein, S. and Zanghì, N. [1992]: ‘Quantum Equilibrium and the Origin of Absolute Uncertainty’, *Journal of Statistical Physics*, **67**, pp. 843–907.
- Dürr, D., Goldstein, S. and Zanghì, N. [2012]: *Quantum Physics without Quantum Philosophy*, Heidelberg: Springer.

- Earman, J. [2006]: ‘The “Past Hypothesis”: Not Even False’, *Studies in History and Philosophy of Modern Physics*, **37**, pp. 399–430.
- Ehrenfest, P. and Ehrenfest, T. [2002]: *The Conceptual Foundations of the Statistical Approach in Mechanics*, Chelmsford, MA: Courier Corporation.
- Esfeld, M. [2014]: ‘Quantum Humeanism, or: Physicalism without Properties’, *The Philosophical Quarterly*, **64**, pp. 453–70.
- Esfeld, M. and Deckert, D.-A. [2017]: *A Minimalist Ontology of the Natural World*, London: Routledge.
- Feynman, R. [2017]: *The Character of Physical Law*, Cambridge, MA: MIT Press.
- Feynman, R. P., Leighton, R. B. and Sands, M. [2015]: *The Feynman Lectures on Physics*, Vol. 1, New York: Basic Books.
- Forrest, P. [1988]: *Quantum Metaphysics*, Oxford: Blackwell.
- Goldstein, S., Lebowitz, J. L., Mastrodonato, C., Tumulka, R. and Zanghi, N. [2010]: ‘Approach to Thermal Equilibrium of Macroscopic Quantum Systems’, *Physical Review E*, **81**, p. 011109.
- Goldstein, S. and Tumulka, R. [2011]: ‘Approach to Thermal Equilibrium of Macroscopic Quantum Systems’, in P. L. Garrido, J. Marro and F. de los Santos (eds), *Non-equilibrium Statistical Physics Today*, New York: American Institute of Physics, pp. 155–63.
- Goldstein, S., Tumulka, R. and Zanghi, N. [2016]: ‘Is the Hypothesis about a Low Entropy Initial State of the Universe Necessary for Explaining the Arrow of Time?’, *Physical Review D*, **94**, p. 023520.
- Goldstein, S. and Zanghi, N. [2013]: ‘Reality and the Role of the Wave Function in Quantum Theory’, in D. Z. Albert and A. Ney (eds), *The Wave Function: Essays on the Metaphysics of Quantum Mechanics*, New York: Oxford University Press, pp. 91–109.
- Hubert, M. and Romano, D. [2018]: ‘The Wave-Function as a Multi-field’, *European Journal for Philosophy of Science*, **8**, pp. 521–37.
- Lebowitz, J. L. [2008]: ‘Time’s Arrow and Boltzmann’s Entropy’, *Scholarpedia*, **3**, pp. 3448.
- Loewer, B. [1996]: ‘Humean Supervenience’, *Philosophical Topics*, **24**, pp. 101–27.
- Loewer, B. [2007]: ‘Counterfactuals and the Second Law’, in H. Price and R. Corry (eds), *Causation, Physics, and the Constitution of Reality: Russell’s Republic Revisited*, Oxford: Oxford University Press.
- Loewer, B. [unpublished]: ‘The Mentaculus Vision’.
- Maroney, O. [2005]: ‘The Density Matrix in the de Broglie–Bohm Approach’, *Foundations of Physics*, **35**, pp. 493–510.
- Maudlin, T. [2007]: *The Metaphysics within Physics*, New York: Oxford University Press.
- Maudlin, T. [2013]: ‘The Nature of the Quantum State’, in D. Z. Albert and A. Ney (eds), *The Wave Function: Essays on the Metaphysics of Quantum Mechanics*, New York: Oxford University Press, pp. 126–53.
- Miller, E. [2014]: ‘Quantum Entanglement, Bohmian Mechanics, and Humean Supervenience’, *Australasian Journal of Philosophy*, **92**, pp. 567–83.
- Myrvold, W. [2017]: ‘Philosophical Issues in Quantum Theory’, in E. N. Zalta (ed.), *The Stanford Encyclopedia of Philosophy*, available at <plato.stanford.edu/archives/spr2017/entries/qt-issues/>.
- Ney, A. [2012]: ‘The Status of Our Ordinary Three Dimensions in a Quantum Universe’, *Noûs*, **46**, pp. 525–60.

- North, J. [2013]: 'The Structure of a Quantum World', in D. Z. Albert and A. Ney (eds), *The Wave Function: Essays in the Metaphysics of Quantum Mechanics*, Oxford: Oxford University Press.
- Penrose, R. [1979]: 'Singularities and Time-Asymmetry', in S. Hawking and W. Israel (eds), *General Relativity*, Cambridge: Cambridge University Press, pp. 581–638.
- Penrose, R. [1999]: *The Emperor's New Mind: Concerning Computers, Minds, and the Laws of Physics*, Oxford: Oxford University Press.
- Price, H. [1996]: *Time's Arrow and Archimedes' Point: New Directions for the Physics of Time*, New York: Oxford University Press.
- Price, H. [2004]: 'On the Origins of the Arrow of Time: Why There is Still a Puzzle about the Low-Entropy Past', in C. Hitchcock (ed.), *Contemporary Debates in the Philosophy of Science*, Oxford: Blackwell, pp. 219–32.
- Sklar, L. [1995]: *Physics and Chance: Philosophical Issues in the Foundations of Statistical Mechanics*, Cambridge: Cambridge University Press.
- Von Neumann, J. [1955]: *Mathematical Foundations of Quantum Mechanics*, Princeton, NJ: Princeton University Press.
- Wallace, D. [2012]: *The Emergent Multiverse: Quantum Theory According to the Everett Interpretation*, Oxford: Oxford University Press.
- Wallace, D. [forthcoming]: 'Probability and Irreversibility in Modern Statistical Mechanics: Classical and Quantum', in D. Bedingham, O. Maroney and C. Timpson (eds), *Quantum Foundations of Statistical Mechanics*, Oxford: Oxford University Press.
- Wallace, D. [unpublished]: 'The Logic of the Past Hypothesis'.
- Wallace, D. and Timpson, C. G. [2010]: 'Quantum Mechanics on Spacetime, I: Spacetime State Realism', *British Journal for the Philosophy of Science*, **61**, pp. 697–727.